

UNCLASSIFIED

AD 400 583

DEFENSE DOCUMENTATION CENTER

FOR

SCIENTIFIC AND TECHNICAL INFORMATION

CAMERON STATION, ALEXANDRIA, VIRGINIA



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

400583

S/830/62/000/002/002/002
D214/D308

STEP

AUTHORS:

Yagodin, G.A. and Chekmarev, A.M.

TITLE:

The extraction of zirconium and hafnium by tri-*n*-octylamine from metal fluoride solutions

SOURCE:

Ekstraktsiya; teoriya, primeneniye, apparatura, no. 2, Ed. by A.P. Zefirov and M.M. Senyavin. Moscow, Gosatomizdat, 1962, 141 - 153

TEXT:

The extraction of Zr and Hf from K_2MF_6 (where M = Zr, Hf) by a solution of pure tri-*n*-octylamine (TOA) in benzene is discussed. TOA will extract Zr and Hf only from weak acid solutions since more acid solutions tend to form $R_3N.HX$ (where X = HSO_4 , Cl, NO_3). Highest values for the distribution coefficients, D_{Zr} and D_{Hf} , were obtained with 0.2 M H_2SO_4 while HNO_3 leads to the lowest D_M values. With H_2SO_4 ,
Card 1/2

The extraction of zirconium ...

S/830/62/000/002/002/002
D214/D308

$D_{Hf} > D_{Zr}$ but with HNO_3 Zr is preferentially extracted. The influence of additions was also studied. The values of D_M decrease as the concentration of the addition in the aqueous phase increases. With small additions of KCl or KF $D_{Hf} > D_{Zr}$ but at higher concentrations ($> 8g/l$ for KCl - $> 1\%$ for KF) $D_{Zr} > D_{Hf}$. Addition of K_2SO_4 make $D_{Zr} > D_{Hf}$ but with NH_4NO_3 $D_{Hf} > D_{Zr}$. Extraction from K_2ZrF_6 (10 g/l) acidified with 0.2 M $(COOH)_2$ by 5% benzene solution of TOA gives $D_{Zr} = 47$ and $D_{Hf} = 10$. Both D values decrease as the molarity of the acid is decreased. The extraction mechanism is summarized by: $2(R_3NH)HSO_{4org} + K_2ZrF_{6aq} \rightleftharpoons (R_3NH)_2ZrF_{6org} + 2KHSO_{4aq}$. Evidence for this mechanism is discussed in detail. There are 12 figures and 6 tables.

Card 2/2